



Welcome and Introduction

R.C. Fernow
BNL
Muon Collider Simulation Workshop
Miami Beach, FL

12 December 2004



Muon Collider Simulation milestones



- first collider concepts ~1969-1971 Budker, Skrinsky
- theory of ionization cooling developed ~1965-1983 Kolomensky, Skrinsky, Parkhomchuk, Balbekov, Neuffer
- early studies followed evolution of transport equations
- early simulations with studied isolated elements ~1994 e.g. Parmela, Simucool
- first Monte Carlo systems studies ~1994 MCM, mix of simple simulation & scaling laws
- development of more flexible codes ICOOL, DPGeant ~1996
- muon collider status report (1999)



Muon collider status report



- most complete overall description of a muon collider ever
- simulations only of isolated parts
- used non-optimal linear cooling channels
- not self-consistent
- emittance exchange was crude



Developments since 1999



- (1) better linear cooling channels developed for neutrino factory studies 1, 2, 2a
- (2) ring coolers
 6D cooling factors ~100
 injection is still major issue
- (3) Balbekov front end design (2003) PR + BCR + RC + linear Li channel
- (4) theory for gas-filled, helical channel → Muons, Inc. early simulation work
- (5) progress on non-scaling FFAGs current NF designs up to 20 GeV need studies for 60 GeV, 1 TeV



Why different from neutrino factory studies?



- (1) lots more cooling required $\sim 10^6$ versus ~ 4 for study 2a
- (2) lots more acceleration required ~1 TeV versus 20 GeV in study 2a
- (3) all particles consolidated in a few bunches 2 versus 80 in study 2a
- (4) need collider ring more difficult than storage ring
- (5) need integrated detector in ring more difficult than isolated detector shielding more critical



Goals for this workshop



- rekindle momentum on muon collider systems simulations
- reexamine critical issues that need to be addressed
- focus ring cooler studies towards specific parts of collider system
- make plans for detailed end-to-end simulation of a muon collider same level of detail as study 2a self-consistent \rightarrow good/bad beam correlations try to reach status report requirements for front end 0.16 μ/p with $\epsilon_{6N}=0.17$ mm³



Humility is a virtue



• we do not have a defendable scenario for a muon collider now

• we should keep an open mind about different approaches

single bunch bunch train with recombination frictional cooling optical stochastic cooling ???

